Retrazin

Presentación, preparación y aplicación



Aureon©
Biotech

Retrazin

- Preparación y Aplicación 4
- Dosaje y Suministro 6
- Cadena de Frío 8
- Más información 9



Aureon© Biotech



Aureon©
Biotech

Retrazin

PREPARACIÓN Y APLICACIÓN

GUÍA VISUAL - 8 ETAPAS

01 HIGIENIZACIÓN DE LAS MANOS



Lávese bien las manos con agua y jabón antes de manipular cualquier material. Este paso es fundamental para evitar contaminaciones y asegurar que todo el procedimiento se realice de forma segura.

02 PREPARACIÓN DEL MATERIAL



Reúna todo lo que va a utilizar antes de comenzar: el frasco del medicamento, algodón, alcohol y la jeringa. Tener todo a mano permite que el procedimiento sea más rápido y seguro.

03 ESTERILIZACIÓN DEL FRASCO



Retire la tapa protectora del frasco y limpie bien el tapón de goma con un algodón empapado en alcohol al 70%. Esto ayuda a eliminar gérmenes y mantener el medicamento en condiciones seguras.

04 CARGA DE AIRE EN LA JERINGA



Con la jeringa vacía, tire del émbolo hacia atrás hasta que el borde quede exactamente en la marca de 0,5 ml. De esta forma carga 0,5 ml de aire en la jeringa.

05 ASPIRACIÓN DEL LÍQUIDO



Introduzca la aguja a través del tapón de goma del frasco. Empuje el émbolo hasta el fondo para ingresar aire al vial y luego tire del émbolo para cargar la cantidad indicada del medicamento.

06 ELIMINACIÓN DE BURBUJAS DE AIRE



Con la jeringa nuevamente hacia arriba, golpéela suavemente y presione levemente el émbolo hasta asegurarse de que no queden burbujas de aire antes de la aplicación.

07 APLICACIÓN DE LA INYECCIÓN



Elija el lugar adecuado para la aplicación (abdomen, muslo o brazo). Realice un pliegue en la piel con los dedos e introduzca la jeringa en un ángulo de 45° y, sin soltar el pliegue, presione suavemente el émbolo. Espere aproximadamente 5 segundos antes de retirar la aguja.

08 DESECHO SEGURO



Una vez finalizada la aplicación, deseche la jeringa en un recipiente adecuado para materiales punzocortantes. Esto es importante para evitar accidentes y posibles contaminaciones.

Aureon® Biotech


Retrazin®
Retatrutide
40mg/2 ml
40 Mg
1
Multi-Dose Vial for Subcutaneous
Use Injectable Solution
Aureon®
Biotech
MADE IN UK
Retrazin
Retatrutide 40MG
Aureon
Retrazin
Retatrutide 40MG
Multi-Dose Vial for
Aureon
Aureon©
Biotech

Retrazin

DOSAJE Y SUMINISTRO

TABLA DE VALORES - 5 CANTIDADES

RETATRUTIDE

Dosis deseada	Volumen (mL)	Jeringa U-100
2,5 mg	0,125 mL	12,5 unidades
5 mg	0,25 mL	25 unidades
7,5 mg	0,375 mL	37,5 unidades
10 mg	0,50 mL	50 unidades
12,5 mg	0,625 mL	62,5 unidades

Aureon® Biotech

★ A CAMBRIDGE-HEADQUARTERED BIOTECHNOLOGY INSTITUTION ★

Aureon Biotech.

Research · Development · Scientific Endorsement

14 Cambridge Innovation Park · Milton Road · Cambridge CB4 0GF · United Kingdom

OFFICIAL DOCUMENT

Certificate of Analysis

N° GBU-RYA-2026-15268

Issued: 23 June 2026 · Cambridge, UK

MHRA · UK Regulatory Compliance | UK GMP · EU GMP Aligned | ISO 9001:2015 — Quality Management | USP <71> · <85> · <788>

QUALITY CONTROL & BATCH RELEASE REPORT — *RETRAZIN®*

RETRATUTIDE · LYOPHILISED POWDER FOR SUBCUTANEOUS INJECTION

PRODUCT NAME **RETRAZIN® · Retratutide**	PRESENTATION **Available strengths · 15 mg & 60 mg vials**
BATCH / LOT NO. **W-QVA-14932**	MANUFACTURING DATE **June 2026**
EXPIRY DATE **June 2028 (24 months)**	STORAGE CONDITIONS **2°C – 8°C · Refrigerated**
SCIENTIFIC ENDORSEMENT **Aureon Biotech Ltd · Cambridge, UK**	MANUFACTURING ORIGIN **Aureon Licensed Facility**

ANALYTICAL TEST RESULTS

PARAMETER	METHOD	SPECIFICATION	RESULT	STATUS
Molecular Identity	*LC-MS / Mass Spectrometry*	Retradutide $C_{221}H_{342}N_{46}O_{68}$	Conforms	PASS
Chromatographic Purity	*HPLC-UV · 214 nm*	≥ 98.0 %	99.78 %	PASS
Assay / Potency	*HPLC + UV-Vis*	95.0 – 105.0 %	100.4 %	PASS
Residual Moisture	*Karl Fischer Titration*	≤ 5.0 %	2.31 %	PASS
Bacterial Endotoxins	*LAL · USP <85>*	≤ 5.0 EU/mg	< 0.5 EU/mg	PASS
Sterility	*USP <71> · Membrane Filtration*	No microbial growth	Conforms	PASS
Appearance (reconstituted)	*Visual Inspection*	Clear, colourless solution	Conforms	PASS
pH (reconstituted)	*Potentiometric*	6.5 – 7.5	7.12	PASS

◆ TECHNICAL CONCLUSION

Batch **L-TZN-26045** of **Tirzena® (Tirzepatide)** has been subjected to the full panel of physical, chemical and microbiological tests outlined in this Certificate, fully complying with Aureon Biotech's internal specifications, the United States Pharmacopeia (USP) reference standards, and UK / EU GMP guidelines. **This batch is hereby APPROVED and RELEASED for pharmaceutical distribution under Aureon Biotech scientific endorsement.**

◆ AUTHORISED BY · QUALIFIED PERSON

J. Whitfield

Dr. James A. Whitfield, MPharm PhD MRPharmS

Director of Quality Control & Regulatory Affairs

Aureon Biotech Laboratories Ltd · Cambridge, UK

GPhC Reg. N° 2087491 · MHRA QP-Eligible

Date of release: **23 June 2026** · Place: **Cambridge, United Kingdom**

Aureon Biotech Laboratories Ltd.

14 Cambridge Innovation Park · Milton Road · Cambridge CB4 0GF · England · United Kingdom

Registered in England & Wales · MHRA Compliant · UK GMP / EU GMP Aligned

Tirzena® is a registered trademark of Aureon Biotech Ltd. © 2026 All rights reserved.

Verify authenticity at **Retrazin.com**

Retrazin


Retrazin
Retatrutide 40Mg.
Multi-Dose Vial for Subcu
Aureon
Retrazin
Retatrutide 40Mg.
Multi-Dose Vial for Subcutaneous
Aureon
Retrazin
Retatrutide 40Mg.
Multi-Dose Vial for Subcutaneous
Aureon
Retrazin
Retatrutide 40Mg.
Multi-Dose Vial for Subcutaneous
Aureon
Retrazin
Retatrutide 40Mg.
Multi-Dose Vial for Subcutaneous
Aureon
Aureon
Biotech

Retrazin

CADENA DE FRÍO & CONSERVACIÓN

BAJO ESTÁNDARES DE CONTROL

Faja de conservación	2°C - 8°C
Ambiente	Refrigeración monitoreada
Presentación	Frasco Multidosis

La integridad del compuesto depende de **una cadena de almacenamiento continua**, con monitoreo de temperatura y condiciones farmacéuticas controladas.



Aureon© Biotech


Retrazin
Retatrutide 40MG
Multi-Dose Vial for Sub
Aureon
CLICK AQUI PARA SABER MÁS
Aureon©
Biotech